



LP LEGACY® PREMIUM SUB-FLOOR ADHESIVE

PRODUCT BENEFITS

- Gun-applied, fast-curing PU collapsible foam adhesive
- 100% solvent-free
- For interior and exterior use
- Time- and money-saving application
- One 29 oz. can = 12X the coverage of a typical tube of conventional sub-floor adhesive
- Protects against squeaks by expanding to create a tight seal in small gaps between panels and supports
- Can be applied to wet, dry or frozen lumber
- Meets ASTM D3498
- VOC compliant
- Excellent adhesion to OSB, treated lumber, masonry, decking, concrete, metals, plywood, etc.
- Squeak-free Lifetime* Limited Warranty when used with LP Legacy® Premium Sub-Flooring Assemblies or LP® TopNotch® 350 Series Sub-Flooring Assemblies

*For complete warranty details, visit LPCorp.com/warranties



Visit LPCorp.com/Legacy or call 1-888-820-0325 today!

PERFORMANCE CATEGORY	PRODUCT PERFORMANCE
Base	Single Component Polyurethane
Consistency	Collapsible Gel Adhesive
Curing System	Moisture Cure at Room Temperature
Curing Time*	About 4 Hours
Skin Formation*	About 20 Minutes
Full Strength*	About 24 Hours
Post Expansion	Minimal
Shear Strength (Wet Lumber) (ASTM D3498)	209 PSI
Shear Strength (Dry Lumber) (ASTM D3498)	422 PSI
Shear Strength (Frozen Lumber) (ASTM D3498)	302 PSI
Moisture Resistance (ASTM D3498)	163 PSI
Temperature Resistance	-40° F to +194° F (+248° F Up to Max. 1 Hour)

*Measured at 68° F/65% R.H. These values may vary depending on ambient factors such as temperature, humidity and type of substrate.

LIMITATIONS

- Excellent adhesion on most substrates, except for PTFE, PE and PP.
- Lower temperature and humidity will extend curing time.



Visit LPCorp.com/Legacy or call 1-888-820-0325 today!

⚠ CAUTION: KEEP OUT OF REACH OF CHILDREN. For professional use only. KEEP AWAY FROM HEAT (>50 °C/122 °F), SPARKS AND OPEN FLAME. DO NOT PUNCTURE OR INCINERATE CAN. DO NOT EXPOSE TO DIRECT SUNLIGHT. USE ONLY IN WELL VENTILATED AREAS. In case of eye contact, flush eye with water for 15 minutes and get immediate medical attention. For inhalation, move to fresh air. For skin contact, wash thoroughly with water and soap. If swallowed, drink large quantities of water, but do not induce vomiting. Call a physician immediately. Refer to SDS for further information.